

# Simultaneous visualization of different genomes (J, JSt and St) in a *Thinopyrum intermedium* × *Thinopyrum ponticum* synthetic hybrid (Poaceae) and in its parental species by multicolour genomic in situ hybridization (mcGISH)

Klaudia Kruppa<sup>1</sup>, Márta Molnár-Láng<sup>1</sup>

<sup>1</sup> Agricultural Institute, Centre for Agricultural Research, Hungarian Academy of Sciences, Department of Plant Genetic Resources, H-2462 Martonvásár, Brunszvik u. 2, Hungary

Corresponding author: Márta Molnár-Láng ([molnar.marta@agrar.mta.hu](mailto:molnar.marta@agrar.mta.hu))

---

Academic editor: L.G. Souza | Received 24 November 2015 | Accepted 27 April 2016 | Published 17 June 2016

---

<http://zoobank.org/A0483A15-477D-455A-A353-C935668FCC60>

---

**Citation:** Kruppa K, Molnár-Láng M (2016) Simultaneous visualization of different genomes (J, JSt and St) in a *Thinopyrum intermedium* × *Thinopyrum ponticum* synthetic hybrid (Poaceae) and in its parental species by multicolour genomic in situ hybridization (mcGISH). Comparative Cytogenetics 10(2): 283–293. doi: 10.3897/CompCytogen.v10i2.7305

---

## Abstract

Multicolour genomic *in situ* hybridization (mcGISH) using total genomic DNA probes from *Thinopyrum bessarabicum* (Săvulescu & Rayss, 1923) Á. Löve, 1984 (genome J<sup>b</sup> or E<sup>b</sup>, 2n = 14), and *Pseudoroegneria spicata* (Pursh, 1814) Á. Löve, 1980 (genome St, 2n = 14) was used to characterize the mitotic metaphase chromosomes of a synthetic hybrid of *Thinopyrum intermedium* (Host, 1805) Barkworth & D.R. Dewey, 1985 and *Thinopyrum ponticum* (Podpěra, 1902) Z.-W. Liu et R.-C. Wang, 1993 named „Agropyron glael” and produced by N.V. Tsitsin in the former Soviet Union. The mcGISH pattern of this synthetic hybrid was compared to its parental wheatgrass species. Hexaploid *Th. intermedium* contained 19 J, 9 J<sup>St</sup> and 14 St chromosomes. The three analysed *Th. ponticum* accessions had different chromosome compositions: 43 J + 27 J<sup>St</sup> (PI531737), 40 J + 30 J<sup>St</sup> (VIR-44486) and 38 J + 32 J<sup>St</sup> (D-3494). The synthetic hybrid carried 18 J, 28 J<sup>St</sup> and 8 St chromosomes, including one pair of J-St translocation and/or decreased fluorescent intensity, resulting in unique hybridization patterns. Wheat line Mv9kr1 was crossed with the *Thinopyrum intermedium* × *Thinopyrum ponticum* synthetic hybrid in Hungary in order to transfer its advantageous agronomic traits (leaf rust and yellow rust resistance) into wheat. The chromosome composition of a wheat/A.glael F<sub>1</sub> hybrid was 21 wheat + 28 wheatgrass (11 J + 14 J<sup>St</sup> + 3 S). In the present study, mcGISH involving the simultaneous use of St and J genomic DNA as probes provided information about the type of *Thinopyrum* chromosomes in a *Th. intermedium*/ *Th. ponticum* synthetic hybrid called A. glael.



**Keywords**

multicolour GISH, *Thinopyrum intermedium*, *Thinopyrum ponticum*, *Agropyron glael*, J, J<sup>st</sup>, St genomes

**Introduction**

N.V. Tsitsin produced a synthetic hybrid in the former Soviet Union by crossing *Thinopyrum intermedium* (Host, 1805) Barkworth & D.R. Dewey, 1985 (= *Agropyron glaucum* Roemer & Schultes, 1817,  $2n=6x=42$ ) with *Thinopyrum ponticum* (Podpěra, 1902) Z.-W.Liu & R.-C.Wang, 1993 (= *Agropyron elongatum* Host ex P. Beauvois, 1812,  $2n=10x=70$ ) (Tsitsin 1954). The hybrid plants were named “*Agropyron glael*” (*A. glael*,  $2n=8x=56$ , Tsitsin 1979), from an abbreviation of “glaucum” and “elongatum”. This name (*A. glael*) will be used hereafter in this article. A number of *A. glael* plants were maintained in Martonvásár (Hungary) thanks to cooperation between the Hungarian Academy of Sciences and the Moscow Research Institute of Agriculture - “Nemchinovka” in the 1960’s. The hybrid plants had 56 chromosomes.

Both wheatgrass species are long been known to have superior resistance to various diseases (Wang 2011). They can be crossed with wheat, making them a potential source of gene pool for wheat improvement. In 2001, wheat line Mv9kr1 was crossed with *A. glael* in Hungary in order to transfer its advantageous agronomic traits (leaf rust and yellow rust resistance) into wheat (Molnár-Láng et al. 2012).

Polyploid *Thinopyrum* (Á. Löve, 1980) species contain genomes similar to the J (E<sup>b</sup>, J<sup>b</sup>) genome of the diploid *Th. bessarabicum* (Săvulescu & Rayss, 1923) Á. Löve, 1984 ( $2n=2x=14$ ) (Östergren 1940) or the E (E<sup>c</sup>, J<sup>c</sup>) genome of *Th. elongatum* (Host, 1802) D.R. Dewey, 1984 ( $2n=2x=14$ ) (Cauderon and Saigne 1961), which are closely related (Ceoloni et al. 2014), and sometimes also contain a third genome (S or St) from *Pseudoroegneria spicata* (Pursh, 1814) Á. Löve, 1980 ( $2n=2x=14$ ). The S genome of *Pseudoroegneria* (Nevski, 1934) genus was renamed to St in order to discriminate from the S genome of Sitopsis section of *Aegilops* Linnaeus, 1753 species (Wang et al. 1995). Ceoloni (2014) also mentioned this genome as St/S. *Th. intermedium* has been described using various genome formulas, including E<sup>c</sup>E<sup>b</sup>St (Wang and Zhang 1996), E<sup>1</sup>E<sup>2</sup>St (Zhang et al. 1996) and JJ<sup>s</sup>S (Chen et al. 1998). Wang et al. (2011) mentioned J<sup>s</sup> as E<sup>St</sup> (J<sup>St</sup>). J<sup>St</sup> symbol will be used hereafter to describe this special chromosome type of *Th. intermedium*. Kishii et al. (2005) and Mahelka et al. (2011) revealed new aspects of its genomic composition, suggesting the possible presence of a *Dasypyrum* (Cosson & Durieu de Maisonneuve, 1855) T. Durand, 1888 (V) genome. Recently Wang et al. (2015) published genotypic data obtained using EST-SSR primers derived from the putative progenitor diploid species *Ps. spicata*, *Th. bessarabicum* and *Th. elongatum*, which indicated that the V genome was not one of the three genomes in intermediate wheatgrass. They proposed the J<sup>vs</sup>J<sup>r</sup>St genome designation, where J<sup>vs</sup> and J<sup>r</sup> represented ancestral genomes of the present-day J<sup>b</sup> of *Th. bessarabicum* and J<sup>c</sup> of *Th. elongatum*, J<sup>vs</sup> being the more ancient. The change of J<sup>s</sup> to J<sup>vs</sup> is based on the study of Mahelka et al. (2011) and Deng et al. (2013), as all 14 chromosomes of J<sup>s</sup> showed GISH/FISH



hybridization signals from V-genome probes [*Dasypyrum villosum* (Linnaeus, 1753) P. Candargy, 1901], but only 8 to 11 of the 14 chromosomes have the centromeric region being hybridized by the St genome probe (Chen et al. 1998, Zhang et al. 1996, Kishii et al. 2005, Tang et al. 2011, Deng et al. 2013). FISH analysis using pMD232-500 as probe (originating from *Secale cereale* Linnaeus, 1753 cv. Kustro) indicated that the 14 J chromosomes of *Th. intermedium* bear FISH signals. According to their findings the J genome is changed to J<sup>r</sup>. The genome constitution of *Th. ponticum* was described using the JJJJ<sup>s</sup>J<sup>s</sup> (Chen et al. 1998) and E<sup>c</sup>E<sup>b</sup>E<sup>x</sup>StSt (Li and Zhang 2002) formulas.

Genomic *in situ* hybridization (GISH) or multicolour genomic *in situ* hybridization (mcGISH) offered new opportunities for testing genome relationships in plants (Bennett et al. 1991), for describing hybrid character (Keller et al. 1996), for visualizing genomes simultaneously (Mukai et al. 1993), and for studying genome organization and evolution (Chen et al. 1994, Mahelka et al. 2011).

Multicolour genomic *in situ* hybridization was used in the present study for the simultaneous visualization of the J and St genomic DNA of *A. glael* and their parental wheatgrass species (*Th. intermedium*, *Th. ponticum*) and to describe the chromosome composition of these materials. As previously published papers had different findings and the authors proposed different genome formulas in *Th. intermedium*, difficulties in identification of the different genomes were expected in our study. As *Th. ponticum* chromosomes belonged to two different genomes (J and J<sup>st</sup>), precise detection and identification of them was probable despite of the high chromosome number. There were no former molecular cytogenetic data about the *A. glael*, but the presence of all the three different chromosome types (J, J<sup>st</sup>, St) of the two parental wheatgrass species was hoped-for.

## Methods

*Thinopyrum intermedium*, *Th. ponticum*, their synthetic hybrid *A. glael*, and the wheat/*A. glael* F<sub>1</sub> hybrid were analysed cytogenetically (Table 1). Seeds of *A. glael*, wheat/*A. glael* F<sub>1</sub> hybrid, *Th. intermedium*, and *Th. ponticum* were germinated, after which mitotic metaphase chromosome spreads were prepared according to Lukaszewski et al. (2004). McGISH was performed in order to simultaneously visualize the different wheatgrass chromosomes in *Th. intermedium*, *Th. ponticum*, *A. glael*, and in the Mv9kr1/*A. glael* F<sub>1</sub> hybrid. J (E<sup>b</sup>) genomic DNA from *Th. bessarabicum* labelled with biotin-11-dUTP (Roche Diagnostics, Mannheim, Germany) and St genomic DNA from *Ps. spicata* labelled with digoxigenin-11-dUTP were produced using the random primed labelling protocol. The hybridization mixture contained 100 ng each of the labelled probes/slide, dissolved in a 15 µl mixture of 100% formamide, 20×SSC and 10% dextran-sulphate at a ratio of 5:1:4, and 3000 ng *Triticum aestivum* (Linnaeus, 1753) DNA (genotype Mv9kr1, BBAADD) as a block when needed. Hybridization was performed at 42°C overnight. Streptavidin-FITC (Roche) and Anti-Digoxigenin-Rhodamine (Roche) dissolved in TNB (Tris-NaCl-blocking buffer) were used in the detection phase. The



**Table 1.** Species and genotypes analysed in the present study.

Genotype	Accession number	Genebank	Geographic origin
<i>Thinopyrum intermedium</i>	PI565004	USDA ARS GRIN	Russia
<i>Thinopyrum ponticum</i>	PI 636523	USDA ARS GRIN	Argentina
<i>Th. ponticum</i>	PI531737	USDA ARS GRIN	Argentina
<i>Th. ponticum</i>	PI 547313	USDA ARS GRIN	Russia
<i>Th. intermedium</i> × <i>Th. ponticum</i> synthetic hybrid: Agropyron glael	glael-8/2008	Martonvásár Cereal Genebank	Russia
Mv9kr1 × A. glael F <sub>1</sub> hybrid	112705	Martonvásár Cereal Genebank	Hungary

slides were screened using a Zeiss Axioskop-2 fluorescence microscope equipped with filter sets appropriate for DAPI (Zeiss Filterset 01), and for the simultaneous detection of FITC and Rhodamine (Zeiss filter set 24). Images were captured with a Spot CCD camera (Diagnostic Instruments) and processed with Image Pro Plus software (Media Cybernetics).

**Results**


*Thinopyrum intermedium*

McGISH, performed using J and St genomic DNA probes, simultaneously discriminated three different genomes in the segmental autoallohexaploid *Th. intermedium* (Fig. 1a–b). Among the 42 chromosomes, 14 fluoresced bright red along their whole length, showing the presence of the St genome. The St probe gave also a hybridization signal in the centromeric region of 9 chromosomes, where the other parts hybridized with the J genome, resulting in two-coloured chromosomes with a bow-tie shape. These J<sup>St</sup>-type chromosomes differed to those of the J, where the chromosomes hybridized with the J genomic DNA probe over the entire length with no centromeric St signals. The intensity of the green fluorescence signal was not uniform, the J<sup>St</sup> chromosomes being fainter than J. In the telomeric segment of some J and J<sup>St</sup> chromosomes a weak St genomic hybridization signal was detected, although in a few other chromosomes this fragment was unlabelled. One satellited chromosome was observed where the NOR region was hybridized to St genomic DNA. The analysed accession (No. PI565004) contained 19 J, 9 J<sup>St</sup> and 14 St chromosomes.


*Thinopyrum ponticum*

The analysed *Th. ponticum* contained 70 chromosomes and two groups could be distinguished based on their mcGISH pattern (Fig. 2). Bright green fluorescence signals





**Figure 1.** Results of multicolour genomic *in situ* hybridization on *Thinopyrum intermedium*. **a** Karyotype of a complete cell using *Thinopyrum bessarabicum* (J, green) and *Pseudoroegneria spicata* (St, red) genomic DNA as probes. Chromosome with satellite is indicated with arrow **b** Karyogram of *Thinopyrum intermedium* chromosomes. Top row: J chromosomes; middle row: J<sup>St</sup> chromosomes with the St pericentromeric region; bottom row: St chromosomes. Bar = 10  $\mu$ m.



**Figure 2.** Multicolour genomic *in situ* hybridization on *Thinopyrum ponticum*. **a** Karyotype of *Th. ponticum* (accession VIR-44486) carrying 40 J and 30 J<sup>St</sup> chromosomes, using *Thinopyrum bessarabicum* (J, green) and *Pseudoroegneria spicata* (St, red) genomic DNA as probes **b** 38 J and 32 J<sup>St</sup> chromosomes identified in *Th. ponticum* (accession D-3494) **c** J<sup>St</sup> chromosomes with different lengths of St DNA in the centromeric region. J<sup>St</sup> chromosomes were marked with asterisks. Bar = 10  $\mu$ m.



marked the J chromosomes, while those with St (red) pericentromeric regions belonged to the J<sup>St</sup> genome. The three analysed accessions showed different chromosome compositions: 43 J + 27 J<sup>St</sup> (PI531737), 40 J + 30 J<sup>St</sup> (PI 547313, Fig. 2a) and 38 J + 32 J<sup>St</sup> (PI636523, Fig. 2b). The length of the St segment in the J<sup>St</sup> chromosome varied (Fig. 2c). Each J<sup>St</sup> chromosome showed a short section of St hybridization close to the centromere, while others fluoresced bright red on almost 1/3 of the chromosomes in the centromeric-pericentromeric regions. There was variation in the intensity of the green fluorescence signal, J<sup>St</sup> chromosomes being fainter than J. The telomeric region of most of the chromosomes did not hybridize with the J or St genomic DNA probes and remained unlabelled.


### ***Thinopyrum intermedium* × *Th. ponticum* synthetic hybrid: *A. glael***

McGISH made it possible to discriminate three different groups of *A. glael* chromosomes (Fig. 3a). The designation of the *A. glael* chromosomes was J, J<sup>St</sup> and St, as the synthetic hybrid contains chromosomes from both *Th. intermedium* and *Th. ponticum*. Digoxigenin-labelled St genomic DNA hybridized to four pairs of submetacentric chromosomes, which were thus identified belonging to the St genome. One pair of chromosomes was mainly red, but an St-J translocation was detected in the long arm (marked with yellow arrowheads). Nine pair of chromosomes with only green fluorescence signals were identified as J genome, though three pairs showed lower fluorescence intensity, while the others were bright. The remaining fourteen pairs had various lengths of St genomic hybridization in the pericentromeric region, showing the presence of the J<sup>St</sup> genome.

### **Wheat/*A. glael* F<sub>1</sub> hybrid**

Chromosome counting detected 49 chromosomes in the wheat/*A. glael* F<sub>1</sub> hybrid (21 wheat + 28 wheatgrass), 28 of which hybridized with the J and/or St genomes during mcGISH, discriminating the wheatgrass chromosomes from the unlabelled wheat (Fig. 3b). Only three chromosomes hybridized with the St genomic DNA over their entire length. Eleven chromosomes had no red fluorescence signal in the centromeric region, and were thus identified as J. Two of them had only very weak J signals, with stronger St hybridization in the telomeric region. The remaining 14 chromosomes had various lengths of St genomic hybridization in the pericentromeric region, showing the presence of the J<sup>St</sup> genome. Some of the J and J<sup>St</sup> chromosomes had a weak St genomic hybridization signal in the telomeric region. Some of these chromosomes carried several J-St, J<sup>St</sup>-St, translocations and/or decreased fluorescent intensity was observed in the pericentromeric and telomeric regions, resulting in unique hybridization patterns (marked with asterisks in Fig. 3c). The chromosome composition of the F<sub>1</sub> hybrid was 21 wheat + 11 J + 14 J<sup>St</sup> + 3 S.





**Figure 3.** Multicolor genomic *in situ* hybridization pattern of *Agropyron glael* and the wheat (Mv9kr1 genotype)/*A. glael* F<sub>1</sub> hybrid. **a** Karyotype of a partial cell of *A. glael* using *Thinopyrum bessarabicum* (J, green) and *Pseudoroegneria spicata* (St, red) DNA probes. Translocation between J and St chromosomes were marked with arrows **b** Karyotype of a complete cell of wheat (Mv9kr1 genotype)/*A. glael* F<sub>1</sub> hybrid using *Th. bessarabicum* (J, green) and *Ps. spicata* (St, red) genomic DNA simultaneously as probes and wheat genomic DNA as block simultaneously **c** Karyogram of *A. glael* chromosomes present in the wheat/*A. glael* F<sub>1</sub> hybrid. Nine *A. glael* chromosomes with hybridization patterns different to their parental species are marked with asterisks. Bar = 10 μm.

## Discussion

GISH or mcGISH, a modification of fluorescence *in situ* hybridization, has been used to characterize genomes and chromosomes in polyploid *Thinopyrum* species (Chen et al. 1998, Tang et al. 2000, Li and Zhang 2002, Mahelka et al. 2011). In the present study, mcGISH involving the simultaneous use of St and J genomic DNA as probes provided information about the number and type of *Thinopyrum* chromosomes and demonstrated the presence of intergenomic (J-St) chromosome rearrangements in *A. glael*.

Chen et al. (1998) used GISH with one labelled genomic DNA probe and one nonlabelled blocking genomic DNA during the characterization of these wheatgrass species. They proposed the symbol J<sup>s</sup> to represent J chromosomes with St repeated sequences and GISH signals around the centromeric regions. This chromosome type was the same which has been described in this study using two labelled genomic DNA probes. The use of mcGISH enabled the J and J<sup>s</sup> genomes of *Th. ponticum* and the J, J<sup>s</sup> and St genomes of *Th. intermedium* to be precisely discriminated using J and St labelled genomic DNA simultaneously.



As the number of J and J<sup>St</sup> chromosomes was usually odd [19 J + 9 J<sup>St</sup> in *Th. intermedium* and 43 J + 27 J<sup>St</sup> (PI531737) in *Th. ponticum*], it is possible that J-J<sup>St</sup> chromosome pairing can occur in meiosis, as reported by Chen et al. (2001). Most of the wheatgrass chromosomes were typical *Thinopyrum* chromosomes in *A. glael* and in the wheat/*A. glael* F<sub>1</sub> hybrid, while others showed notable differences when the mcGISH patterns were compared to those of *Th. ponticum* and *Th. intermedium*: decreased fluorescence intensity, J-St translocations in the telomeric region of J<sup>St</sup> chromosomes, and unlabelled chromosome parts in all types of chromosomes. Chen et al. (2001) reported a high frequency of chromosome pairing between J-J<sup>St</sup>, J-St and J<sup>St</sup>-St chromosomes, as the result of which genetic exchange is possible between these genomes. Several minor J-St and J<sup>St</sup>-St translocations were observed in *A. glael* and the wheat/*A. glael* F<sub>1</sub> hybrid. These translocations may have occurred during the formation of the synthetic hybrids. As the J-J<sup>St</sup>-St chromosomes paired at high frequency, it may be that *A. glael* is not only a hybrid of the two wheatgrass species, but that the genetic composition has changed or been enriched with DNA sequences from other species during the long maintenance period (decades), as wheatgrass species are open-pollinating and very polymorphic. This could explain the presence of different hybridization patterns between the wheatgrass chromosomes in *A. glael* and the wheat/*A. glael* F<sub>1</sub> hybrid.

Several types of genome composition and chromosome numbers have been reported for *Th. intermedium* (Chen et al. 1998, Tang et al. 2000, Da Yong et al. 2004). Chen et al. (1998) detect 41 chromosomes (18 J, 10 J<sup>St</sup>, 13 St) in a line derived from Portugal (PI249145), 49 chromosomes (18 J, 10 J<sup>St</sup>, 21 St) in a French genotype. Most of the analyzed accessions carried 42 chromosomes, but the number of each chromosome type was various: 20 J + 8 J<sup>St</sup> + 14 St in 'Chef', 'Clarke' (USA), 18 J + 10 J<sup>St</sup> + 14 St in PI317406 (Afghanistan), and 21 J + 7 J<sup>St</sup> + 14 St in PI547333 (China) (Chen et al. 1998). Tang et al. (2000) analyzed a Chinese accession and identified 21 J + 7 J<sup>St</sup> + 14 St chromosomes. Da Yong et al. (2004) could detect 28 J + J<sup>St</sup> and 14 St chromosomes in PI 469214 (USA), PI 578698 (Russia), and Z1141 (Canada). In this study we could detect 42 chromosomes including 19 J, 9 J<sup>St</sup> and 14 St. According to other findings and our results, when the chromosome number was not 42, the number of St chromosomes was derived. The number of J + J<sup>St</sup> chromosomes was always 28. Mahelka et al. (2011) detected 42 chromosomes in different *Th. intermedium* accessions, and 14 of which hybridized with *Dasyphyrum villosum* genomic DNA, and also carried St genomic DNA hybridization signal in the pericentromeric region. Mahelka et al. (2011) concluded that the genomic heterogeneity of intermediate wheatgrass was higher than had been assumed, making this species more interesting as a source of desirable agronomic traits.

Nucleolar dominance, an epigenetic phenomenon in which one parental set of ribosomal RNA (rRNA) genes is silenced in an interspecific hybrid or during allopolyploidization, first reported in the 1930s (Navashin 1934). Only ribosomal RNA genes inherited from one parent are transcribed (Pikaard 2000), and the nucleolus organiser regions (NORs), the sites of rRNA genes from the other parent(s) are suppressed. The phenomenon was observed in several interspecific hybrids (Gautam et al. 2014), including wheat/*Thinopyrum elongatum* addition lines (Linc et al. 2012). *Thinopyrum intermedium* and *Thinopyrum ponticum* are allopolyploid species, thus nucleolar dominance



can probably be observed in them, and especially in their synthetic hybrid (A.glael). Loss of secondary constrictions can be observed during allopolyploidization or the formation of interspecific hybrids, which can be studied using rDNA probes by FISH. It was not part of this study, but it is planned in the future. Allopolyploidy can induce rapid genome evolution, and can cause genomic shock. The nature of this phenomenon were investigated (Matsuoka 2011). Its manifestation includes chromosomal rearrangement, the gain and loss of chromosome segments, gene repression and activation, subfunctionalization, transposon activation, and changes in the epigenome (Wang et al. 2014). Multicolour GISH is a powerful technique to detect interspecific and intergeneric chromosome rearrangement. According to our mcGISH results, we observed minor chromosomal rearrangement, St-J translocations in nine chromosomes of A.glael in the wheat/ A.glael F1 hybrid, which chromosome patterns couldn't observed in *Thinopyrum* parental species. The reduction of number of St chromosomes were also detected.

As A. glael contains chromosomes from the two most valuable *Thinopyrum* species, changes in its genome could result in new invaluable genetic material, especially for wheat breeding.

## Conclusions

In the present study, mcGISH involving the simultaneous use of St and J genomic DNA as probes provided information about the genome composition and the type of *Thinopyrum* chromosomes in a *Th. intermedium*/ *Th. ponticum* synthetic hybrid called A. glael.

## Acknowledgements

This work was funded by the Hungarian National Scientific Research Fund (OTKA K 104382 and K 108555). Special thanks to Dezső Szalay, and to the Moscow Research Institute of Agriculture - "Nemchinovka", who kindly provided the A. glael plants. The authors gratefully acknowledge the excellent technical assistance of F. Tóth. Thanks are due to Barbara Hooper for revising the manuscript linguistically.

## References

- Bennett ST, Kenton AY, Bennett MD (1992) Genomic *in situ* hybridization reveals the allopolyploid nature of *Milium montianum* (Gramineae). *Chromosoma* 101(7): 420–424. doi: 10.1007/BF00582836
- Chen Q, Armstrong K (1994) Genomic *in situ* hybridization in *Avena sativa*. *Genome* 37(4): 607–612. doi: 10.1139/g94-086
- Chen Q, Conner RL, Laroche A, Thomas JB (1998) Genome analysis of *Thinopyrum intermedium* and *Thinopyrum ponticum* using genomic *in situ* hybridization. *Genome* 41(4): 580–586. doi: 10.1139/gen-41-4-580



- Chen Q, Conner RL, Laroche A, Ahmad F (2001) Molecular cytogenetic evidence for a high level of chromosome pairing among different genomes in *Triticum aestivum* – *Thinopyrum intermedium* hybrids. *Theoretical and Applied Genetics* 102(6): 847–852. doi: 10.1007/s001220000496
- Cauderon Y, Saigne B (1961) New interspecific and intergeneric hybrids involving *Agropyrum*. *Wheat Info Service* 12: 13–14.
- Ceoloni C, Kuzmanovic L, Gennaro A, Forte P, Giorgi D, Grossi MR, Bitti A (2014) Genomes, chromosomes and genes of the wheatgrass genus *Thinopyrum*: the value of their transfer into wheat for gains in cytogenomic knowledge and sustainable breeding. In: Tuberosa R, Graner A, Frison E (Eds) *Genomics of Plant Genetic Resources, Crop Productivity, Food Security and Nutritional Quality Vol. 2*, Springer Science+Business Media, Dordrecht, 333–358. doi: 10.1007/978-94-007-7575-6\_14
- Da-Yong L, Yan-Yan R, Xue-Yong Z (2004) Chromosomal distribution of the 18S-5.8S-26S rDNA loci and heterogeneity of nuclear ITS regions in *Thinopyrum intermedium* (Poaceae: Triticeae). *Acta Botanica Sinica* 46(10): 1234–1241.
- Deng CL, Bai LL, Fu SL, Yin WB, Zhang YX, Chen YH, Wang RRC, Zhang XQ, Han FP, Hu ZM (2013) Microdissection and chromosome painting of the alien chromosome in an addition line of wheat-*Thinopyrum intermedium*. *PLoS ONE* 8(8): e72564. doi: 10.1371/journal.pone.0072564
- Gautam M, Ge XH, Li ZY (2014) Brassica. In: Pratao A, Kumar J (Eds) *Alien Gene Transfer in Crop Plants, Volume 2, Achievements and Impacts*, Springer Science+Business Media, 207–230. doi: 10.1007/978-1-4614-9572-7\_10
- Keller ERJ, Schubert I, Fuchs J, Meister A (1996) Interspecific crosses of onion with distant *Allium* species and characterization of the presumed hybrids by means of flow cytometry, karyotype analysis and genomic in situ hybridization. *Theoretical and Applied Genetics* 92(3): 417–424. doi: 10.1007/BF00223688
- Kishii M, Wang RRC, Tsujimoto H (2005) GISH analysis revealed new aspect of genomic constitution of *Thinopyrum intermedium*. *Proceedings of the 5th International Triticeae Symposium, Prague, June 6–10, 2005, Czech Journal of Genetics and Plant Breeding* 41: 91–95.
- Li D, Zhang X (2002) Physical localization of the 18S-5.8S-26S rDNA and sequence analysis of ITS regions in *Thinopyrum ponticum* (Poaceae: Triticeae): Implications for concerted evolution. *Annals of Botany* 90(4): 445–452. doi: 10.1093/aob/mcf213
- Linc G, Sepsi A, Molnár-Láng M (2012) A FISH karyotype to study chromosome polymorphisms for the *Elytrigia elongata* E genome. *Cytogenet Genome Research* 13: 138–144. doi: 10.1159/000334835
- Lukaszewski AJ, Rybka K, Korzun V, Malyshev SV, Lapinski B, Whitkus R (2004) Genetic and physical mapping of homoeologous recombination points involving wheat chromosome 2B and rye chromosome 2R. *Genome* 47(1): 36–45. doi: 10.1139/g03-089
- Mahelka V, Kopecký D, Paštová L (2011) On the genome constitution and evolution of intermediate wheatgrass (*Thinopyrum intermedium*: Poaceae, Triticeae). *BMC Evolutionary Biology* 11: 127. doi: 10.1186/1471-2148-11-127
- Matsuoka Y (2011) Evolution of polyploid *Triticum* wheats under cultivation: the role of domestication, natural hybridization and allopolyploid speciation in their diversification. *Plant and Cell Physiology* 52(5): 750–764. doi: 10.1093/pcp/pcr018



- Molnár-Láng M, Linc G, Szakács É, Molnár I, Cseh A, Schneider A, Kruppa K (2012) Wheat-alien introgression programme in Martonvásár. III Vavilov International Conference, Russian Academy of Sciences, St. Petersburg, November 6–9: 239–240.
- Mukai Y, Nakahara Y, Yamamoto M (1993) Simultaneous discrimination of the three genomes in hexaploid wheat by multicolor fluorescence *in situ* hybridization using total genomic and highly repeated DNA probes. *Genome* 36(3): 489–494. doi: 10.1139/g93-067
- Navashin N (1934) Chromosome Alterations Caused by Hybridization and Their Bearing upon Certain General Genetic Problems. *Cytologia* 5(2): 169–203. doi: 10.1508/cytologia.5.169
- Östergren G (1940) Cytology of *Agropyron junceum* and *A. repens* and their spontaneous hybrids. *Hereditas* 26(3-4): 305–316. doi: 10.1111/j.1601-5223.1940.tb03239.x
- Pikaard CS (2000) The epigenetic of nucleolar dominance. *Trends in Genetics* 16(11): 495–500.
- Tang S, Li Z, Jia X, Larkin PJ (2000) Genomic *in situ* hybridization (GISH) analyses of *Thinopyrum intermedium*, its partial amphiploid Zhong 5, and disease-resistant derivatives in wheat. *Theoretical and Applied Genetics* 100(3): 344–352. doi: 10.1007/s001220050045
- Tang ZX, Yang ZJ, Fu SL, Yang MY, Li GR, Zhang HQ, Tan FQ, Ren Z (2011) A new long terminal repeat (LTR) sequence allows the identify J genome from Js and St genomes of *Thinopyrum intermedium*. *Journal of Applied Genetics* 52(1): 31–33. doi: 10.1007/s13353-010-0019-8
- Tsitsin NV (1979) Cytogenetic studies of wheatgrasses and wheat-wheatgrass hybrids. [In Russian: Цитогенетические исследования видов пырея и пшенично-пырейных гибридов]. In: Tsitsin NV (Ed.) Problems of distant hybridization. Nauka Press, Moscow, 48–53. [In Russian: Проблемы отдаленной гибридизации]
- Wang H, Jang J, Chen S, Qi X, Fang W, Guan Z, Teng N, Liao Y, Chen F (2014) Rapid Genetic and Epigenetic Alterations under Intergeneric Genomic Shock in Newly Synthesized *Chrysanthemum morifolium* × *Leucanthemum paludosum* Hybrids (Asteraceae). *Genome Biology and Evolution* 6(1): 247–259. doi: 10.1093/gbe/evu008
- Wang RRC (2011) *Agropyron* and *Psathyrostachys*. In: Kole C (Ed.) *Wild Crop Relatives: Genomic and Breeding Resources, Cereals*. Springer, Berlin, 77–108. doi: 10.1007/978-3-642-14228-4\_2
- Wang RRC, von Bothmer R, Dvorak J, Fedak G, Linde-Laursen I, Muramatsu M (1995) Genome symbols in the *Triticeae* (*Poaceae*). In: Wang RRC, Jensen KB, Jaussi C (Eds) *Proceedings of 2nd International Triticeae Symposium*, Utah State University Publications on Design and Production, Logan, 29–34.
- Wang RRC, Zhang XY (1996) Characterization of the translocated chromosome using fluorescence *in situ* hybridization and random amplified polymorphic DNA on two *Triticum aestivum*-*Thinopyrum intermedium* translocation lines resistant to wheat streak mosaic or barley yellow dwarf virus. *Chromosome Research* 4(8): 583–587. doi: 10.1007/BF02261721
- Wang RRC, Larson SL, Jensen KB, Bushman BS, DeHaan LR, Wang S, Yan X (2015) Genome evolution of intermediate wheatgrass as revealed by EST-SSR markers developed from its three progenitor diploid species. *Genome* 58(2): 63–70. doi: 10.1139/gen-2014-0186
- Zhang XY, Koul A, Petroski R, Ouellet T, Fedak G, Dong YS, Wang RRC (1996) Molecular verification and characterization of BYDV resistant germ plasms derived from hybrids of wheat with *Thinopyrum ponticum* and *Th. intermedium*. *Theoretical and Applied Genetics* 93(7): 1033–1039. doi: 10.1007/BF00230121